

Abstract of the Invention

The subject invention provides a compound having the formula:

H1 - X - B - Y - H2

wherein each of H1 and H2 may be the same or different and capable of binding to a receptor which is the same or different; wherein each of X and Y may be present or absent and if present, each may be the same or different spacer moiety; and wherein B is an enzyme cleavable moiety. This invention also provides a method of screening proteins for the ability to catalyze bond cleavage, comprising the steps of:

- a) providing a cell that expresses a pair of fusion proteins which upon dimerization change a cellular readout;
- b) providing the compound of the invention which dimerizes the pair of fusion proteins, said compound comprising two portions coupled by a bond that is cleavable by the protein to be screened; and
- c) screening for the cellular readout, wherein a change the cellular readout indicates catalysis of bond cleavage by the protein to be screened. Finally, the invention also provides a method of screening proteins for the ability to catalyze bond formation, comprising the steps of:
- a) providing a cell that expresses a pair of fusion proteins which upon dimerization activate a cellular readout:
- b) providing a first compound and a second compound, each being capable of binding to one of the pair of fusion proteins, said first and second compound comprising a portion through which the first and second compounds are coupled to form the inventive compound by the action of the bond forming protein to be screened; and
- c) screening for the cellular readout, wherein a change in the cellular readout indicates catalysis of bond formation by the protein to be screened.

